

Ferromagnetic to antiferromagnetic transition of one-dimensional spinor Bose gases with spin-orbit coupling

Xing Chen,¹ Haiping Hu,¹ Yuzhu Jiang,² and Shu Chen^{1,*}

¹*Beijing National Laboratory for Condensed Matter Physics,
Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*

²*Beijing Computational Science Research Center, Beijing 100084, China*
(Dated: February 28, 2013)

We have analytically solved one-dimensional interacting two-component bosonic gases with spin-orbit (SO) coupling by the Bethe-ansatz method. Through a gauge transformation, the effect of SO coupling is incorporated into a spin-dependent twisted boundary condition. Our result shows that the SO coupling can influence the eigenenergy in a periodical pattern. The interplay between interaction and SO coupling may induce the energy level crossing for the ground state, which leads to a transition from the ferromagnetic to antiferromagnetic state.

PACS numbers: 67.85.-d, 67.60.Bc, 03.75.Mn

I. INTRODUCTION

The experimental success in manipulating cold atoms in effective one-dimensional (1D) waveguides has deepened our understanding of the properties of the low-dimensional quantum gases [1–4]. Meanwhile, studies of synthetic gauge field in cold atom systems have also made great progresses: pioneering experiments of NIST group have generated the effective magnetic fields, electric fields and spin-orbit (SO) coupling in ultracold Bose gases [5], and SO-coupled Fermi gases have also been realized recently [6, 7]. Systems with SO coupling have revealed intriguing phenomena in condensed matter physics, such as quantum spin Hall effect and topological insulators [8, 9], where electrons play the elemental role in these physical systems. The realization of SO coupling in cold bosonic systems opens a completely new avenue for studying the interesting physics beyond the traditional condensed matter physics.

Many theoretical researches have investigated and predicted interesting phenomena for SO coupled bosonic systems. For example, a single plane wave phase or a stripe phase in spin-1/2 Bose-Einstein condensates (BECs) with SO coupling has been predicted depending on the intraspecies interaction larger or smaller than that of interspecies [10, 11], collective modes [12] and stability [13, 14] of BECs with SO coupling have been discussed, and interesting phases in the presence of harmonic traps and rotation have also been studied [15–20]. As most of these researches are restricted on mean-field approximation in the weakly interacting regime, there is a rather basic question in what extent these results are applicable for an arbitrary interaction. This makes the exact solution for SO coupled cold atom system highly desirable.

In this paper, we analytically solve the SO coupled spin-1/2 bosonic gases in a ring trap by the Bethe-

ansatz (BA) method and find that the SO coupling can bring some new physics to the system particularly in the strongly interacting regime. In the absence of SO coupling, the spin-1/2 bosonic system is solved analytically by the BA method and the ground state is found to be the ferromagnetic state [21–24]. However, this result is largely changed by adding the SO coupling. Our results based on solving BA equations indicate that the SO coupling influences the eigenenergy periodically. In the strongly interacting regime, we find that the interplay of the SO coupling and interaction may produce level crossing for the ground state. Consequently, the system may change from a ferromagnetic ground state to an antiferromagnetic state.

The present paper is organized as follows. Section II introduces the model. By using a rather general transformation, SO coupling effect can be transformed to the twisted boundary condition. The exact solution is given by the BA method and the energy level crossing is shown in the energy spectrum. Section III discusses the BA equations in the strongly interacting limit. The antiferromagnetic state is proved to be the ground state in the infinite interaction limit. A summary is given in Section IV.

II. MODEL AND SOLUTION

We consider a two-component bosonic gas confined in a 1D ring trap in the presence of SO coupling with the Hamiltonian given by $\hat{H} = \hat{H}_0 + \hat{H}_{\text{int}}$ with

$$\hat{H}_0 = \int dx \hat{\Psi}^\dagger(x) \left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} - 2i\alpha\sigma_z \frac{\partial}{\partial x} \right) \hat{\Psi}(x), \quad (1)$$

where $\hat{\Psi}^\dagger = (\hat{\Psi}_\uparrow^\dagger, \hat{\Psi}_\downarrow^\dagger)$ represents two internal states of bosonic atoms, α denotes the spin-orbit coupling strength and σ_z is the Pauli matrix. Here, we consider a quasi-one-dimensional situation with the transverse motion tightly confined in its ground state. The ring trap enforces the periodic boundary condition $\hat{\Psi}_\sigma(x) = \hat{\Psi}_\sigma(x + L)$.

*Electronic address: schen@aphy.iphy.ac.cn

The interaction term is generally represented as $\hat{H}_{\text{int}} = \int dx (g_1 \hat{n}_\uparrow^2 + g_2 \hat{n}_\downarrow^2 + 2g_{12} \hat{n}_\uparrow \hat{n}_\downarrow)$, where $g_{1,2}$ and g_{12} denote the strengths of intraspecies and interspecies interaction, which are experimentally tunable. In this work, we shall focus on the case with spin-independent interaction, i.e., $g_1 = g_2 = g_{12} = c$, for which interaction term can be represented as $\hat{H}_{\text{int}} = c \int dx \hat{n}^2$ with $\hat{n} = \hat{n}_\uparrow + \hat{n}_\downarrow$. We shall set $\frac{\hbar^2}{2m} = 1$ in the following text for convenience.

It is difficult to solve the Hamiltonian with spin and momentum coupled together. Using a rather general transformation [25]

$$\hat{\Psi}_\sigma(x) = \hat{\Phi}_\sigma(x) e^{-i\alpha x \sigma_z}, \quad (2)$$

the Hamiltonian is rewritten as follows,

$$\hat{H}_0 = - \sum_\sigma \int dx \hat{\Phi}_\sigma^\dagger(x) \frac{\partial^2}{\partial x^2} \hat{\Phi}_\sigma(x) - N\alpha^2, \quad (3)$$

and the form of $\hat{H}_{\text{int}} = c \int dx \hat{n}^2$ is invariant with $\hat{n} = \hat{\Phi}^\dagger \hat{\Phi}$ and $\hat{\Phi}^\dagger = (\hat{\Phi}_\uparrow^\dagger, \hat{\Phi}_\downarrow^\dagger)$. The operator $\hat{\Phi}_\sigma(x)$, $\hat{\Phi}_\sigma^\dagger(x')$ also satisfy commutation relation $[\hat{\Phi}_\sigma^\dagger(x), \hat{\Phi}_{\sigma'}(x')] = \delta_{\sigma\sigma'} \delta(x-x')$. Meanwhile, the total momentum now is represented as $\hat{K} = \sum_\sigma \int dx \hat{\Phi}_\sigma^\dagger(x) (-i\frac{\partial}{\partial x} - \alpha\sigma_z) \hat{\Phi}_\sigma(x)$ which is spin-dependent. The Schrödinger equation is $\hat{H}|\Phi\rangle = E|\Phi\rangle$ with wave function

$$|\Phi\rangle = \sum_\kappa \int \phi_\kappa(x_1 \dots x_N) \prod_{i=1 \dots N} \hat{\Phi}_{\sigma_i}^\dagger(x_i) dx_i |0\rangle, \quad (4)$$

here κ denotes $\sigma_1, \sigma_2, \dots, \sigma_N$ corresponding to the spin index for different particles. By applying the wave function to the Schrödinger equation, we get

$$H\phi_\kappa(x_1, \dots, x_N) = E\phi_\kappa(x_1, \dots, x_N) \quad (5)$$

with

$$H = - \sum_{i=1}^N \frac{\partial^2}{\partial x_i^2} + 2c \sum_{i < j} \delta(x_i - x_j) - \alpha^2 N, \quad (6)$$

It is worth mentioning that SO effect is not vanishing but transformed to the spin-dependent twist boundary condition. Explicitly, the periodic boundary condition $\hat{\Psi}_\sigma(x+L) = \hat{\Psi}_\sigma(x)$ under the transformation Eq.(2) is changed as

$$\hat{\Phi}_\sigma(x+L) = \hat{\Phi}_\sigma(x) e^{i\sigma_z \alpha L}, \quad (7)$$

Correspondingly, the wave function now should satisfy

$$\phi_\kappa(x_1, \dots, x_j + L, \dots, x_N) = e^{i\sigma_{j,z} \alpha L} \phi_\kappa(x_1, \dots, x_j, \dots, x_N). \quad (8)$$

The model of two-component bosons described by Eq.(6) is analytically solvable by Bethe ansatz (BA) under periodic boundary conditions [21]. The eigenstates can be characterized by the total spin S of the system

which varies from 0 to $N/2$. The ground state of interacting two-component bosons corresponds to the ferromagnetic state with $S = N/2$. Now the problem of solving one-dimensional interacting two-component bosonic gases with SO coupling reduces to solve the integrable model of (6) under the twist boundary condition of (8), for which we can still obtain exact solutions by the same method as originally developed in [26]. The system is solvable by the same Bethe-type wavefunction given by

$$\begin{aligned} \phi_\kappa(x_1 \dots x_N) &= \sum_{P,Q} \theta(x_{q_N} - x_{q_{N-1}}) \dots \theta(x_{q_2} - x_{q_1}) \\ &\times A(Q, P) e^{i \sum_j k_{p_j} x_{q_j}}, \end{aligned} \quad (9)$$

where $Q = (q_1, q_2, \dots, q_N)$ and $P = (p_1, p_2, \dots, p_N)$ denote permutations of $1, \dots, N$, $A(Q, P)$ are coefficients to be determined, $\theta(x_2 - x_1)$ is the step function and $\{k_j\}$ are quasimomenta. The coefficients fulfill the following relations

$$A(Q; \dots i, j \dots) = Y_{ji}^{ab} A(Q; \dots j, i \dots), \quad (10)$$

with $Y_{jl}^{ab} = \frac{(k_j - k_l) P_{q_a q_b} - ic}{k_j - k_l + ic}$, P_{q_a, q_b} permutes q_a, q_b in $A(\dots q_a, q_b, \dots, P)$. The scattering matrix remains the same as in the model with periodic boundary conditions. For eigenstate with total spin $S = \frac{1}{2}(N - 2M)$ ($0 \leq M \leq N/2$) under the twist boundary condition of (8), we obtain the following Bethe-ansatz equations

$$e^{i(k_j - \alpha)L} = - \prod_{l=1}^N \frac{k_j - k_l + ic}{k_j - k_l - ic} \prod_{\beta=1}^M \frac{k_j - \lambda_\beta - ic'}{k_j - \lambda_\beta + ic'}, \quad (11)$$

$$\prod_{j=1}^N \frac{\lambda_\zeta - k_j - ic'}{\lambda_\zeta - k_j + ic'} = -e^{i2\alpha L} \prod_{\beta=1}^M \frac{\lambda_\zeta - \lambda_\beta - ic}{\lambda_\zeta - \lambda_\beta + ic}, \quad (12)$$

with $c' = c/2$, $\{k_j\}$ the quasimomenta and $\{\lambda_\beta\}$ the spin rapidities. From the BA equations, one can observe that k depends on α periodically. So we can take $\alpha = \alpha_0 + \frac{2\pi n}{L}$ with $\alpha_0 \in [0, \frac{2\pi}{L}]$ and n an integer. In fact, we need only consider $\alpha_0 \in [0, \frac{\pi}{L}]$. The quasimomentum for $\alpha_0 \in [\frac{\pi}{L}, \frac{2\pi}{L}]$ can be deduced from $\alpha_0 \in [0, \frac{\pi}{L}]$ by two steps: first taking $-\alpha_0 \in [-\frac{\pi}{L}, 0]$ with solution $-k_j$ and $-\lambda_\zeta$, secondly taking $-\alpha_0 + \frac{2\pi}{L} \in [\frac{\pi}{L}, \frac{2\pi}{L}]$ with the solution unchangeable.

Taking logarithm of the above equations, we get

$$\begin{aligned} (k_j - \alpha)L &= 2\pi I_j - \sum_{l=1}^N 2 \tan^{-1} \frac{k_j - k_l}{c} \\ &+ \sum_{\beta=1}^M 2 \tan^{-1} \frac{k_j - \lambda_\beta}{c'}, \end{aligned} \quad (13)$$

$$\sum_{j=1}^N 2 \tan^{-1} \frac{\lambda_\zeta - k_j}{c'} = 2\pi J_\zeta + 2\alpha L + \sum_{\beta} 2 \tan^{-1} \frac{\lambda_\zeta - \lambda_\beta}{c}, \quad (14)$$




FIG. 1: (Color online) The lowest energy spectrum versus c for systems with $N = 16$, $\alpha = 0$ and different total spins. The ground state is a ferromagnetic state and the energy levels satisfy $E(S) < E(S')$ for $S > S'$.

where I_j with $j = 1, \dots, N$ denote the density quantum numbers and J_ζ with $\zeta = 1, \dots, M$ the spin quantum numbers. Here I_j and J_ζ are integer (half-integer) depending on $N - M$ is odd (even). Taking the period of α into account, let $I_j = I_{j_0} - n$ and $J_\zeta = J_{\zeta_0} - 2n$, the above equations are reduced to

$$(k_j - \alpha_0)L = 2\pi I_{j_0} - \sum_{l=1}^N 2 \tan^{-1} \frac{k_j - k_l}{c} + \sum_{\zeta=1}^M 2 \tan^{-1} \frac{k_j - \lambda_\zeta}{c'}, \quad (15)$$

$$\sum_{j=1}^N 2 \tan^{-1} \frac{\lambda_\zeta - k_j}{c'} = 2\pi J_{\zeta_0} + 2\alpha_0 L + \sum_{\beta=1}^M 2 \tan^{-1} \frac{\lambda_\zeta - \lambda_\beta}{c}. \quad (16)$$

The corresponding eigenenergy is given by

$$E = \varepsilon - N\alpha^2 \quad (17)$$

with $\varepsilon = \sum_j k_j^2$. The total momentum is given by

$$K = \tilde{K} - 2S\alpha, \quad (18)$$

with

$$\tilde{K} = \sum_j k_j = \frac{2\pi}{L} \left(\sum_{j=1}^N I_{j_0} - \sum_{\zeta=1}^M J_{\zeta_0} \right) + 2S\alpha_0. \quad (19)$$




FIG. 2: (Color online) The lowest energy spectrum versus c for systems with $N = 16$, $\alpha = 0.95k_c/N$ and different total spins. The level crossing is shown and the ground state is the antiferromagnetic state in the strong interaction limit.

In the absence of the SO coupling, the model reduces to the exactly solvable interacting two-component bosonic model [21–23] with the ground state being a ferromagnetic state with $S = N/2$, which is degenerate for $S_z \in [-S, S]$. For a given S , we can calculate the lowest eigenenergy by solving the BA equations. To give a concrete example, we calculate the system with $N = 16$ and show the lowest energy in a given total spin space versus the interaction strength in Fig.1 for different total spin S . Here the parameters c and E are in units of k_c and Nk_c^2 , respectively, with k_c defined as $k_c = \pi\rho = \pi N/L$. Apparently, we have $E(S) < E(S')$ for $S > S'$ in the whole regime of interaction strength, and the ground state is a ferromagnetic state with maximum S . This is in contrast to the case of spin-1/2 fermionic gases, for which Lieb-Mattis theorem [27] gives the opposite law: $E(S) < E(S')$ for $S < S'$ and the ground state is an antiferromagnetic state with $S = 0$. As the interaction tends to the strong interaction limit, the energies for different S approach to the same value and become degenerate in the infinite interaction limit [28–30].

In the presence of the SO coupling, the ground state energy spectrum for systems with different S can be also calculated by solving the corresponding BA equations, which is shown in Fig. 2 as a function of the interaction strength for a fixed SO coupling $\alpha = 0.95k_c/N = 0.95\pi/L$. In contrast to Fig.1, the energy spectrum is largely changed. For weak interaction, the energies still satisfy the rule of $E(S) < E(S')$ for $S > S'$. However, with the increase of interaction, level crossing appears and the ferromagnetic state is no longer the ground state. For large enough interaction, the energy levels fulfill the relation of $E(S) < E(S')$ for $S < S'$. That is to say, the ground state of the two-component bosonic system can change from the ferromagnetic state to the antiferromag-

netic state due to the interplay between the SO coupling and interaction.

III. STRONG COUPLING LIMIT

To see more clearly how the SO coupling affects the ground state energy, we discuss the strong interaction limit, which permits us to get some analytical expressions for the energy spectrum. In the limit of $L|c| \gg 1$, λ_ζ are proportional to the interaction strength c whereas k_j remain finite [31]. Applying the Taylor expansion to Eq.(15) and Eq.(16), the equations of k_j and λ_ζ are simplified as

$$\left(k_j - \frac{\tilde{K}}{N}\right) = \frac{2\pi}{L}I_{j_0} - \frac{2\pi}{LN} \sum_{j=1}^N I_{j_0} - \frac{2\rho}{c} \left(k_j - \frac{\tilde{K}}{N}\right) \left(1 - \frac{v}{2N}\right), \quad (20)$$

$$2N \tan^{-1} \frac{\lambda_\zeta}{c'} = 2\pi J_{\zeta_0} + 2\alpha_0 L + \sum_{\beta=1}^M 2 \tan^{-1} \frac{\lambda_\zeta - \lambda_\beta}{c} \quad (21)$$

with $v = \sum_{\zeta=1}^M \frac{1}{1/4 + (\lambda_\zeta/c)^2}$. For the state of $M = 0$ (or even M), $I_{j_0} = -(N-1)/2 \dots (N-1)/2$, from Eq.(20)

$$\tilde{k}_j = \frac{2\pi}{L}I_{j_0} - \frac{2\rho}{c}\tilde{k}_j \left(1 - \frac{v}{2N}\right)$$

with $\tilde{k}_j = k_j - \frac{\tilde{K}}{N}$. For elementary spin excitations of $M = 1$ (odd M) [22], $I_{j_0}^1 = -N/2, \dots, N/2 - 1$, then

$$\tilde{k}_j = \frac{2\pi}{L} \left(I_{j_0}^1 + \frac{1}{2}\right) - \frac{2\rho}{c}\tilde{k}_j \left(1 - \frac{v}{2N}\right)$$

Since $I_{j_0}^1 + \frac{1}{2} = I_{j_0}$, in general, the equations for quasimomentum k_j are given by

$$k_j = \frac{2\pi I_{j_0}}{L \left(1 + \frac{2\rho}{c} \left(1 - \frac{v}{2N}\right)\right)} + \frac{\tilde{K}}{N}, \quad (22)$$

In the strong coupling limit $c/\rho \rightarrow \infty$, $k_j = 2\pi I_{j_0}/L + \tilde{K}/N$. Without SO coupling, the quasimomenta $\{k_j\}$ are independent on the total spin. From Eq.(19), it can be seen that $\tilde{K}$ is dependent on the total spin by the item $2\alpha_0 S$. In contrast, with SO coupling, the quasimomenta $\{k_j\}$ are shifted by the value of $2\alpha_0 S/N$ for various total spin S . Substituting the above equation into $\varepsilon = \sum_j k_j^2$, we get

$$\varepsilon = \frac{\pi^2}{3} \frac{N(N^2 - 1)}{L^2 \left(1 + \frac{2\rho}{c} \left(1 - \frac{v}{2N}\right)\right)^2} + \frac{\tilde{K}^2}{N}. \quad (23)$$

From Eq.(23), it can be seen that the ground energy depends on SO coupling parameter α from three aspects.




FIG. 3: (Color online) The lowest energy spectrum versus α for systems with $N = 16$, $c/k_c = 1000$ and different total spins. Here E is in units of Nk_c^2 while α is in units of k_c/N . The energy spectrum cyclically changes in a larger scope when $\alpha \in (0, k_c)$.




FIG. 4: (Color online) An enlargement of the energy spectrum shown in Fig.3 for half period of $\alpha \in (0, \pi/L)$ is shown ($\pi/L = k_c/N$). There is energy crossing in this scope. The ferromagnetic state is the ground state while $\alpha = 0$. With the increase of α , the ground state changes to the antiferromagnetic state.

Firstly, the spin rapidity λ_ζ is dependent on α from its self-consistent Eq.(21). Secondly, from Eq.(19), $\tilde{K}$ is proportional to α_0 , and therefore the second item $\tilde{K}^2/N$ is determined by α_0 . Thirdly, the last item $-N\alpha^2$ is a constant for fixed α and N . In the strong interaction limit $c/\rho \rightarrow \infty$, as $\rho/c \rightarrow 0$, the contribution of spin rapidity can be ignored. Only the item $\tilde{K}^2/N$ will affect the energy for different S . Here $\tilde{K}$ is proportional to S from Eq.(19), which leads to the energy ε increasing with the

increase of S . Therefore, SO coupling favors antiferromagnetic state as the ground state.

In the case of even N and M , the quantum numbers $J_{\zeta_0} = -(M+1)/2 - \zeta$ where $\zeta = 1 \dots M$. From Eq.(22), the quasimomenta $k_j = 2\pi I_{j_0}/L + 2\alpha_0 S/N$. From Eq.(23), while $c/\rho \rightarrow \infty$, the energy is given by

$$\frac{\varepsilon}{k_c^2 N} = \frac{N^2 - 1}{3N^2} + \left(\frac{\alpha_0}{k_c}\right)^2 \left(\frac{2S}{N}\right)^2. \quad (24)$$

Apparently, the energy is lowest for $S = 0$. For large but finite interaction, the interplay between SO coupling and interaction leads to the energy level crossing. In Fig. 3, we show the energy spectrums versus α in the strongly interacting case with $c/k_c = 1000$, where the SO coupling parameter is taken in the regime of $\alpha/k_c \in (0, 1)$. The cyclical change of spectrum with the increase of α coincides with our previous discussion. In Fig. 4, the scope of SO coupling parameter α is just half the period $\alpha/k_c \in (0, \pi/L)$ of Fig. 3. With the increase of α , there are energy level crossing for different S . When $\alpha = 0$, the energy levels satisfy $E(S) < E(S')$ for $S < S'$. When α is large enough, the energy levels are opposite to that law. After the crossing, the larger α being, the larger energy difference for various S will emerge.

Before ending the paper, we would like to give a remark on the interacting spin-1/2 fermionic model with SO coupling, for which the SO coupling induced level crossing is absent. As for the 1D interacting spin-1/2 fermionic gas [32], the ground state is an antiferromagnetic state ($S = 0$) in the absence of SO coupling. In the presence of SO coupling, one can still use the gauge transformation to transform the problem into an integrable spin-1/2 fermionic model with a spin-dependent twist boundary condition [25, 26], and the system is determined by the following BA equations

$$(k_j - \alpha) L = 2\pi I_j - \sum_{\zeta=1}^M 2 \tan^{-1} \frac{k_j - \lambda_{\zeta}}{c'}, \quad (25)$$

$$\sum_{j=1}^N \tan^{-1} \frac{\lambda_{\zeta} - k_j}{c'} = 2\pi J_{\zeta} - 2\alpha L + \sum_{\beta=1}^M 2 \tan^{-1} \frac{\lambda_{\zeta} - \lambda_{\beta}}{c}. \quad (26)$$

In the strong interaction limit, the energy is

$$\varepsilon = \frac{\pi^2}{3} \frac{N(N^2 - 1)}{L^2 (1 + \frac{v}{cL})^2} + \frac{\tilde{K}^2}{N}.$$

Similar to the bosonic case, the SO coupling still favors the antiferromagnetic state as the ground state. So the SO coupling does not lead to level crossing for spin-1/2 fermions, i.e., the ground state will always be an antiferromagnetic state.

IV. SUMMARY

In summary, we have analytically solved 1D interacting spin-1/2 bosonic gases with SO coupling. As the effect of SO coupling can be absorbed into the twisted boundary condition, we get the exact solution to this system by BA method and find that the corresponding eigenenergies are periodically dependent on the SO coupling. The interplay between interaction and SO coupling has revealed the existence of energy level crossing and the ground state phase transition from the ferromagnetic state to antiferromagnetic state.

Acknowledgments

We thank Y. J. Hao for helpful discussions. This work has been supported by National Program for Basic Research of MOST, NSF of China under Grants No.11121063 and No.11174360 and 973 grant.

-
- [1] B. Paredes, *et al.*, Nature **429**, 277 (2004).
 - [2] T. Kinoshita, T. Wenger and D. S. Weiss, Science **305**, 1125 (2004).
 - [3] M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac, and M. Rigol, Rev. Mod. Phys. **83**, 1405 (2011).
 - [4] X.-W. Guan, M. T. Batchelor and C. H. Lee, arXiv:1301.6446.
 - [5] Y.-J. Lin *et al.*, Nature(London) **462**, 628 (2009); Y.-J. Lin *et al.*, Nat. Phys. (London) **7**, 531 (2011); Y.-J. Lin, K. J. Garcis, I. B. Spielman, Nature(London) **83**, 471 (2011).
 - [6] P. Wang, Z.-Q. Yu, Z. K. Fu, J. Miao, L. H. Huang, S. J. Chai, H. Zhai, and J. Zhang, Phys. Rev. Lett. **109**, 095301(2012).
 - [7] L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M. W. Zwierlein, Phys. Rev. Lett. **109**, 095302(2012).
 - [8] X. L. Qi and S. C. Zhang, Phys. Rev. Lett. **52**, 2111 (2010).
 - [9] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
 - [10] C. Wang, C. Gao, C.-M. Jian, and H. Zhai, Phys. Rev. Lett. **105**, 160403 (2010).
 - [11] T.-L. Ho and S. Z. Zhang, Phys. Rev. Lett. **107**, 150403 (2011).
 - [12] W. Zheng and Z. Li, Phys. Rev. A **85**, 053607 (2012).
 - [13] T. Ozawa and G. Baym, Phys. Rev. Lett. **109**, 025301 (2012).
 - [14] Q. Z. Zhu, C. W. Zhang, B. Wu, EPL **100**, 50003 (2012).
 - [15] C. J. Wu, I. Mondragon-Shem and X. F. Zhou, Chin. Phys. Lett. **28**, 097102 (2011).
 - [16] B. Ramachandhran B. Opanchuk, X.-J. Liu, H. Pu, P. D. Drummond and Hui Hu, Phys. Rev. A **85**, 023606 (2012).

- [17] S. Sinha, R. Nath, and L. Santos, Phys. Rev. Lett. **107**, 270401 (2011)
- [18] X.-Q. Xu and J. H. Han, Phys. Rev. Lett. **107**, 200401 (2011).
- [19] J. Radić, T. Sedrakyan, I. B. Spielman, and V. Galitski, Phys. Rev. A **84**, 063604 (2011).
- [20] Z. F. Xu, R. Lu, and L. You, Phys. Rev. A **83**, 053602 (2011); Z. F. Xu, Y. Kawaguchi, L. You, and M. Ueda, Phys. Rev. A **86**, 033628 (2012).
- [21] Y. Q. Li, S. J. Gu, Z.J. Ying and U. Echern, Europhys. Lett. **61**, 368 (2003).
- [22] J. N. Fuchs, D. M. Gangardt, T. Keilmann, and G. V. Shlyapnikov, Phys. Rev. Lett. **95**, 150402 (2005).
- [23] X. W. Guan and M. T. Batchelor and M. Takahashi, Phys. Rev. A **76**, 043617 (2007).
- [24] Y. J. Hao, Y. B. Zhang, X. W. Guan and S. Chen, Phys. Rev. A **79**, 033607 (2009).
- [25] S. Uchino and N. Kawakami, Phys. Rev. A **85**, 013610 (2012).
- [26] B. S. Shastry and B. Sutherland, Phys. Rev. Lett. **65**, 243 (1990).
- [27] E. H. Lieb and D. Mattis, Phys. Rev. **125**, 164 (1962).
- [28] M. D. Girardeau and A. Minguzzi, Phys. Rev. Lett. **99**, 230402 (2007).
- [29] L. Guan, S. Chen, Y. Wang, and Z. Q. Ma, Phys. Rev. Lett. **102**, 160402 (2009).
- [30] F. Deuretzbacher, K. Fredenhagen, D. Becker, K. Bongs, K. Sengstock, and D. Pfannkuche, Phys. Rev. Lett. **100**, 160405 (2008).
- [31] L. Guan and S. Chen, Phys. Rev. Lett. **105**, 175301 (2010).
- [32] C. N. Yang, Phys. Rev. Lett. **19**, 1312 (1967).